



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

In vitro studies on odontogenic tumors

Catón, Javier ; Mitsiadis, Thimios A ; Morgan, Peter R

Abstract: Ameloblastomas are uncommon benign neoplasms of the jaws. They originate from dental epithelial cells, but they are not capable of mineralizing or forming enamel. The study of these tumors is limited to live tissue collected from patients during scheduled surgery. Ameloblastomas grow slowly in vivo and this property is translated to their behavior in vitro. Here, we describe the methods to culture ameloblastomas in organotypic cultures, as well as to isolate stem/progenitor cells from these tumors.

DOI: https://doi.org/10.1007/978-1-61779-860-3_15

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-68955>

Journal Article

Originally published at:

Catón, Javier; Mitsiadis, Thimios A; Morgan, Peter R (2012). In vitro studies on odontogenic tumors. *Methods in Molecular Biology*, 887:167-177.

DOI: https://doi.org/10.1007/978-1-61779-860-3_15

Metadata of the chapter that will be visualized online

Series Title	Methods in Molecular Biology	
Chapter Title	In Vitro Studies on Odontogenic Tumors	
Chapter SubTitle		
Copyright Year	2012	
Copyright Holder	Springer Science + Business Media, LLC	
Corresponding Author	Family Name	Catón
	Particle	
	Given Name	Javier
	Suffix	
	Division	Departamento de Anatomía y Embriología Humana I, Faculty of Medicine
	Organization	Universidad Complutense Madrid
	Address	Madrid, Spain
	Email	javicaaton@med.ucm.es
Author	Family Name	Mitsiadis
	Particle	
	Given Name	Thimios A.
	Suffix	
	Division	Department of Orofacial Development and Regeneration, Faculty of Medicine
	Organization	Institute of Oral Biology, ZZM, University of Zurich
	Address	Zurich, Switzerland
	Email	
Author	Family Name	Morgan
	Particle	
	Given Name	Peter R.
	Suffix	
	Division	Oral Pathology
	Organization	King's College London Dental Institute
	Address	London, UK
	Email	
Abstract	Ameloblastomas are uncommon benign neoplasms of the jaws. They originate from dental epithelial cells, but they are not capable of mineralizing or forming enamel. The study of these tumors is limited to live tissue collected from patients during scheduled surgery. Ameloblastomas grow slowly in vivo and this property is translated to their behavior in vitro. Here, we describe the methods to culture ameloblastomas in organotypic cultures, as well as to isolate stem/progenitor cells from these tumors.	
Key words: (separated by '-')	Ameloblastomas - Odontogenic tumors - Enamel - Organotypic culture - Tumor stem cells - Cell cocultures	

In Vitro Studies on Odontogenic Tumors

Javier Catón, Thimios A. Mitsiadis, and Peter R. Morgan

Abstract

Ameloblastomas are uncommon benign neoplasms of the jaws. They originate from dental epithelial cells, but they are not capable of mineralizing or forming enamel. The study of these tumors is limited to live tissue collected from patients during scheduled surgery. Ameloblastomas grow slowly in vivo and this property is translated to their behavior in vitro. Here, we describe the methods to culture ameloblastomas in organotypic cultures, as well as to isolate stem/progenitor cells from these tumors.

Key words: Ameloblastomas, Odontogenic tumors, Enamel, Organotypic culture, Tumor stem cells, Cell cocultures

1. Introduction

Odontogenic tumors (OTs) present considerable challenges for any investigator willing to use cell and organotypic culture in studies with human tissue as the starting material. These challenges could be summarized as follows:

- 1. The range and diversity of the tumors
- 2. The rarity of individual types of odontogenic tumors
- 3. The frequent, although not exclusive, intra-osseous location of these tumors
- 4. The diverse tissue composition of the odontogenic tumors
- 5. The usually slow rate of the odontogenic tumors' growth

If due consideration is paid to these drawbacks, it is possible to employ the tissue in experimental, not simply descriptive, investigations.

The classification of OTs most used currently is based on that published in 2005 by the World Health Organization (WHO) (1), although unfortunately it introduced several somewhat arbitrary changes in terminologies from those in common use. Broadly, the benign OTs are classified along embryological lines according to whether neoplastic odontogenic epithelium appears to reflect interaction with odontogenic ectomesenchyme or not. One subgroup appears to represent neoplastic growth of tissues derived from the ectomesenchyme itself. Malignant OTs are generally classified descriptively according to their similarity to their benign counterparts. [AU1]

Benign OTs represent a range of growth disorders from unequivocal neoplasms (e.g., ameloblastomas) to unequivocal hamartomas (compound and complex odontomas) with some entities having an intermediate status (e.g., adenomatoid odontogenic tumor). Parallels with normal tooth development break down with some tumors because these produce unique structures and/or cells not found in the developing teeth (e.g., ghost cells in the calcifying odontogenic cyst, now termed the calcifying cystic odontogenic tumor). These examples illustrate some of the range and diversity exhibited by OTs.

OTs are uncommon tumors. The most common unequivocal OT is the ameloblastoma, a locally aggressive benign neoplasm. Ameloblastomas represent less than 5% of head and neck neoplasms. In the Afro-Caribbean ethnic group, they are more common and in parts of Africa they represent a significant proportion of untreated neoplasms. Controversially, in 2005, the WHO included a cyst, the odontogenic keratocyst, among odontogenic neoplasms, based upon molecular genetic criteria and its propensity for recurrence. This instantly made this cystic lesion the most common odontogenic neoplasm in ethnic Caucasian and Asians, but the status of this entity is still not settled (2). Adenomatoid odontogenic tumors and the odontomes are the next common OTs, after ameloblastomas, both manifesting in a younger population, the second decade. Ameloblastomas and odontogenic keratocysts peak in the fourth or fifth decades. Most other odontogenic tumors are very rare indeed and, unfortunately for the experimental scientist, usually inaccurately diagnosed preoperatively. The commonest malignant OT is the ameloblastic carcinoma, which is more rare than its benign counterpart. A compendium of incidence data for OTs is to be found in Reichart & Philipsen (3).

Most OTs are intra-osseous, a minority of them arising in the gingiva (i.e., peripheral OTs). This makes for difficulties in accessing tumorigenic tissue for culture or banking fresh frozen samples. Band-saw slicing, appropriate for fixed hard tissues, is contraindicated in fresh specimens for reasons of infection control. However, the behavior of the tumors sometimes assists access. In those OTs that expand the jaws, typically large ameloblastomas, the normally dense cortical bone is thinned and can be prized open after slicing [AU2]




Fig. 1. H&E sections of the two most common solid/multicystic ameloblastomas: Follicular and Plexiform.

with a scalpel to access the soft tumorigenic tissue. Samples taken in this way are less likely to be contaminated with oral microorganisms than those from tumors exposed to the mouth.

A feature of OTs that may pose a problem for diagnosis as well as constituting a disadvantage for cell studies is the extensive cystic change. This is well illustrated in most of the large ameloblastomas. As the neoplasm enlarges, multiple cystic spaces in the epithelial component (i.e., microcysts) and/or the delicate connective tissue (i.e., stromal cysts) expand and coalesce so that opening an expanded cortex reveals a space filled with straw-colored fluid. This is a frequent finding in the two commonest subtypes of ameloblastomas, the solid/multicystic (Fig. 1) and the unicystic variants. Incisional biopsies that include epithelium only from the expanded cyst wall hamper diagnosis, as it is thin and may not show classical features of the tumor and if this is the only material available for cell culture the epithelial cell yield is low.

Other OTs, such as the adenomatoid odontogenic tumor and calcifying odontogenic cyst, may present with a largely cystic expansion of the jaw. The odontogenic keratocyst, assuming we regard it as an OT, has the disadvantage from the perspective of cell culture of not expanding the jaw, or doing so only in juveniles or after a long period of neglect. On the other hand, odontogenic keratocysts are usually treated conservatively nowadays by the delicate detachment of the cyst wall from the inner surface of the jaw (i.e., endosteal), so experimental samples may be obtained direct from the surgeon or pathologist.

Apart from the rare and serendipitous presentation of many OTs, a further problem for cell culture and analytical studies is their heterogeneous nature. Particularly, the rarer OTs contain a mixture of hard and soft tissues. Where the hard tissue is a tooth or several discrete tooth-like elements, these may be removed before explanting or freezing the tissue, but some OTs have dispersed

dentine- or enamel-like hard tissues from which odontogenic epithelium, or even mesenchymal stroma, may be impossible to separate. Two of the most common tumors of this type are mature Pindborg tumors (i.e., calcifying epithelial odontogenic tumors) and the cementoblastoma, one category of OT of odontogenic ectomesenchymal origin consisting almost solely of mineralized tissue.

In the following, we describe methods for the culture of ameloblastoma explants taking into consideration the above difficulties for the collection of this tumor. We do not mention any ethical considerations on the understanding that each researcher will follow the protocols of the institution where the research will take place.

2. Materials

Tissue should arrive at the pathology department fresh. The tissue is divided for pathology, tissue bank, and research (see Note 1).

2.1. Cryopreservation

Liquid nitrogen is commercially available (see Note 2), and stored in liquid nitrogen dewars and liquid nitrogen storage containers. Cryo tubes, Cryo 1°C Freezing Container such as “Mr. Frosty,” and cryo-protectants such as glycerol or dimethyl sulfoxide (DMSO) for cellular cryopreservation.

2.2. Fixatives

Two different types of fixatives are routinely used: 10% neutral buffered formalin for pathological studies and 4% paraformaldehyde in 0.1 M phosphate buffer (PB) for research.

1. 10% Buffered formalin:

Formaldehyde (37–40%)	100 ml
Distilled water	900 ml
NaH ₂ PO ₄	4.0 g
Na ₂ HPO ₄ (anhydrous)	6.5 g

Mix to dissolve. Store at room temperature.

2. 4% Paraformaldehyde in 0.1 M phosphate buffer.

Paraformaldehyde	40 g
0.1 M Phosphate buffer	1,000 ml

Heat to 60–65°C while stirring. Add a few drops of 1 N NaOH until solution is clear. Continue to stir to dissolve. Cool the solution, filter, and aliquot (see Note 3).

2.3. Buffers

1. 0.2 M Phosphate buffer, pH 7.4. 137

Na ₂ HPO ₄	21.8 g	t3.1
NaH ₂ PO ₄	6.4 g	t3.2 138

Distilled water up to 1,000 ml. 139

2. 1 M Tris-Cl (pH 7.4). 140

Tris base	121.1 g	t4.1
HCl, concentrated	70 ml	t4.2 141

Distilled water up to 1,000 ml. 142

3. 10× Tris/EDTA (TE). 143

1 M Tris-Cl	100 ml	t5.1
0.5 M EDTA, pH 8.0	20 ml	t5.2 144

Distilled water up to 1,000 ml (see Note 4). 145

4. HANKs (commercially available for cell culture). 146

5. PBS (commercially available for cell culture). 147

2.4. Tissue Sectioning
and Staining

1. SuperFrost Plus glass slides or similar slides to increase adherence. 148

2. Microtome/cryostat and material related to their use. 149

3. Wax for embedding (purified paraffin/synthetic resin blend). 150

4. Xylene/Histoclear (see Note 5). 151

5. Alcohol gradient, 50, 70, 80, 90, and 100% (see Note 6). 152

6. Tissue-Tek CRYO-OCT Compound, Sucrose, cryo-embed-
ding molds, and -80°C freezer. 153
154

7. Routine H&E staining material. 155

2.5. Culture Media
and Materials

1. Progenitor cell targeted to oral epithelium defined liquid cul-
ture medium (CnT24). 156
157

2. DMEM supplemented with 1× penicillin/streptomycin antibi-
otics for washing and tissue transport. 158
159

3. DMEM supplemented with 10% fetal calf serum (FCS) and 1×
penicillin/streptomycin antibiotics for organotypic cultures. 160
161

4. Differentiation medium: BGJb medium supplemented with
10 mg/ml ascorbic acid, 2 mM of sodium β-glycerophosphate,
and 1× penicillin/streptomycin antibiotics. 162
163
164

5. 2.4 U/ml of Dispase. 165

6. 0.25% Trypsin. 166

7. Cell and organ culture dishes, metal grids, and support filters. 167

2.6. Tools for
Dissection

1. Glass dish. 168

2. Forceps, scalpels, blades, and needles. 169

3. Methods

3.1. Tissue Extraction/
Preservation

Once surgery of the tumor has been scheduled, it is important to follow up with the surgical team. Most hospitals will have a service to bring the tissue to the pathology department (see Note 7), but it is recommended to collect it personally—if possible—to minimize the time from surgery to the laboratory.

3.2. Tissue Selection

A pathologist familiar with ameloblastomas should select an area of the tumor that will be likely to be richer in tumor cells. As mentioned in the introduction, some of these tumors could have very little starting material to work with. The tissue is then divided for pathology, tissue bank (see Note 8), and research. This methods chapter focuses only on the research portion.

3.3. Tissue for Research

The tissue selected for research should be divided:

3.3.1. Fresh Tissue
for In Vivo Studies

1. Flash freeze
This is achieved by submerging the sample in liquid nitrogen or a mixture of dry ice and ethanol. This frozen tissue can be later used to extract nucleic acid for gene expression and genetic studies (qPCR, microarray, etc.).
2. Fixation
Although some tissue will be fixed using 10% buffered formalin (see Note 9), most tissue for research should be fixed immediately by submerging it in 4% PFA at 4°C overnight. The fixed tissue can be used for genetic studies (generally, we perform in situ hybridization, immunohistochemistry).
For in situ hybridization, cryosections are normally used.
3. Cryoprotection
After fixing, the tissue should be rinsed at room temperature in 0.1 M phosphate buffer with 5% sucrose (this process will initiate the cryoprotection of the tissue). Continue with increasing concentrations of sucrose starting at 5% sucrose in phosphate buffer reaching 20% in 5% increments. Proceed for 30 min in each sucrose mixture at room temperature leaving it overnight at 4°C in fresh 20% sucrose/phosphate.
4. Infiltration
The tissue is then placed into an infiltration mixture (2:1, 20% sucrose phosphate buffer and O.C.T. embedding medium) for 30 min at room temperature before freezing (see Note 10).
5. Embedding and freezing
Transfer the tissue to an embedding mold and fill the mold with fresh infiltration mixture. Rapidly submerge the mold into Isopentane cooled with liquid nitrogen (see Note 11). After the material is frozen, wrap the block and store at -80°C.

6. Sectioning	212
3–5- μm sections are cut at -20°C in a cryostat. To achieve	213
ideal sections, it is critical to have the knife-edge as sharp as	214
possible. Trim the block face to a diamond shape, with the	215
long axis oriented vertically. This orientation helps to make	216
removal of the sections from the knife-edge easier, and will	217
minimize handling damage of the tissue. Use a small camel	218
hairbrush to guide the section off the block face and transfer it	219
to glass slides (see Note 12). Allow the section to dry on the	220
slide at room temperature. Store slides at -80°C until needed.	221
 3.3.2. Fresh Tissue for In Vitro Studies	
Fresh tissue can be cultured in an organotypic form to maintain the	222
architectural three dimensions of the tissue or in monolayer cell	223
culture to isolate specific cells.	224
 1. Organotypic cultures	225
Tissue is cut into 2–5-mm cubes and placed into Trowell-type	226
culture dishes (4). These cultures can be maintained for	227
approximately 15 days using DMEM medium supplemented	228
with 10% FCS. The tissue can be induced using proteins and/	229
or cell cocultures. The usage of beads for induction has the	230
advantage of showing the effect in the precise site of applica-	231
tion. Following are examples of induction beads and cell	232
cocultures.	233
 2. Induction beads	234
Affi-gel agarose (75–150- μm diameter) or heparin acrylic	235
beads (100–200 mesh/100–250- μm diameter), depending on	236
the type of proteins, are needed for induction. Ameloblastomas	237
are induced with proteins involved in tooth development.	238
Recombinants are diluted with 0.1% bovine serum albumin	239
(BSA) in PBS, pH 7.4, to concentrations of 100–200 ng/ μl .	240
Recombinant BMP2, BMP4 (Fig. 2), and SHH (200 ng/ μl)	241
are used to preload affi-gel agarose beads and FGF2, FGF3,	242
and FGF4 (100 ng/ μl) are used to preload heparin acrylic	243
beads. These preloaded beads are incubated for 30 min at	244
room temperature and then washed for 5–15 min in culture	245
media before being transferred with a mouth-controlled capillary	246
pipette on top of the explants. As controls, beads loaded with	247
0.1% BSA in PBS are used. The explants with beads are cultured	248
in serum-free medium for 20 h and processed for in situ hybrid-	249
ization (5, 6), proliferation assays (7), and immunohistochemis-	250
try (6, 8) as described in the references (see Note 13).	251
 3. Cell cocultures	252
The usage of cell cocultures for induction of ameloblastoma	253
tissues allows exposing the tumors to a collection of factor the	254
cells express in vitro. We use murine embryonic odontogenic	255
cells (pre-ameloblasts or pre-odontoblasts) to determine the	256
effect of each specific cell type on the explant.	257




Fig. 2. Preloaded affi-gel agarose beads (BMP4 beads) implanted on top of an ameloblastoma (AB) organotypic culture.

Dissected first mandibular molars from E16.5 mouse (see Note 14) are placed in 2.4 U/ml of Dispase and incubated for 1 h at 4°C. The mesenchymal tissue is mechanically separated from the epithelium using tungsten needles. The two tissues are placed separately in 0.25% trypsin at 37°C for 30 min. Disaggregating is achieved by passing the cells after trypsin treatment through an 18-g needle. The cell suspension is centrifuged and the pellet washed in DMEN with 1× pen/strep. [AU3] The cells are centrifuged once more and the pellet is placed using a pipette tip in the ameloblastoma explants (Fig. 3). These explants are cultured in differentiation medium for up to 10 days. The cocultured explants are fixed and treated for study as described before.

4. Monolayer cultures

Cells are isolated from the fresh tissue using two methods:

Explants shedding allows cells to cast off from small tissue explants into a culture dish. The explants are cut in a similar way as the organotypic cultures. These explants are then placed in 10-cm cell culture dishes with CnT24 media just covering the bottom of the dishes. This allows the explants to have an optimum gas exchange. The explants are kept in this culture conditions for approximately 1 week or when epithelial-like cell colonies are observed attaching to the culture dish. The tissue can also be digested in collagenase IV (freshly prepared [AU4]




Fig. 3. GFP expressing odontogenic cells (GFP) placed on top of ameloblastoma (AB) explants in organotypic cocultures.

[AU5]

500 CDU/ml) at 37°C for 30 min. The cells' disaggregation is achieved by passing them through an 18-g needle. The cell suspension is centrifuged and the pellet washed in DMEN with 1× pen/strep. Centrifuge the cells once more and resuspend the pellet in CnT24 medium for plating in culture dishes (see Note 15).

Ameloblastoma cells can then be characterized for markers common with other epithelial stem cells and other tumor stem cells. These markers can be detected by immunohistochemistry, in situ hybridization, or RT-PCR. For membrane-bound markers, the cells can be sorted using fluorescence activated cell sorting (FACS) (9). This allows separating living cells expressing the marker of interest from the rest.

4. Notes

1. Chemicals are purchased commercially from your choice of provider and solutions should be prepared nuclease free for studies with the need for RNA preservation.

2. Liq N2 should be handled with extreme care and protective clothing, gloves, and eye shield should be worn.
3. 4% PFA aliquots can be stored long term at -20°C . Avoid repeated temperature change cycles and bring to near room temperature before use. It is recommended to use phosphate buffer made with nuclease-free water.
4. We also purchase molecular biology-grade 100 \times TE for nucleic acid work.
5. HistoClear is less toxic and the results are similar.
6. Dehydration of tissue for nucleic acid work—in situ hybridization—should be done with gradient alcohols made with nuclease-free water.
7. We normally received the tissue from the pathology department. They select a portion of the tumor and hand it to the research team on DMEM with Pen/Strep. [AU6]
[AU7]
8. Most departments will have a tissue bank for storage of the tissue. This could be useful for in vivo studies.
9. The pathology department that provides the tissue will normally process it in this manner for routine histological analysis. These preparations could be used for morphological study of the tissue.
10. The tissue should sink to the bottom of the container to indicate a correct infiltration.
11. Rapid freezing is recommended, although we have observed that simply placing the mold in dry ice will freeze the sample quickly enough without any adverse effect. It is also possible to use dry ice and ethanol mixture to accelerate the freezing process.
12. If the cryostat has auto-sectioning mode, then one should slow the speed of sectioning (approx. 5 mm/s) to ease the manipulation of the sections as they are coming off the block face. Do not use the anti-roll plate furnished with the cryostat; it compresses the sections and results in poor tissue morphology.
13. In situ hybridization can be done on whole mount or cryosections.
14. We use mice expressing green fluorescent protein (GFP) in order to being able to distinguish the mouse cells from the tumor cells.
15. Epithelial cells are more likely to grow when plated in higher concentrations.

339 **References**

- 340 1. Barnes, L. et al. (2005) *Pathology and Genetics of*
341 *Head and Neck Tumours*, IARC Press, Lyon.
- 342 2. Li, T. J. (2011) The odontogenic keratocyst: a
343 cyst, or a cystic neoplasm? *J Dent Res* 90,
344 133–142.
- 345 3. Reichart, P. A., and Philipsen, H. P. (2003)
346 [Revision of the 1992 edition of the WHO his-
347 tological typing of odontogenic tumors. A sug-
348 gestion], *Mund Kiefer Gesichtschir* 7, 88–93.
- 349 4. Trowell, O. A. (1954) A modified technique
350 for organ culture in vitro, *Exp Cell Res* 6,
351 246–248.
- 352 5. Mitsiadis, T. A., Hirsinger, E., Lendahl, U., and
353 Goridis, C. (1998) Delta-notch signaling in
354 odontogenesis: correlation with cytodifferenti-
355 ation and evidence for feedback regulation, *Dev*
356 *Biol* 204, 420–431.
- 357 6. Mitsiadis, T. A., Salmivirta, M., Muramatsu,
358 T., Muramatsu, H., Rauvala, H., Lehtonen,
359 E., Jalkanen, M., and Thesleff, I. (1995)
360 Expression of the heparin-binding cytokines,
361 midkine (MK) and HB-GAM (pleiotrophin) is
associated with epithelial-mesenchymal inter- 362
actions during fetal development and organo- 363
genesis, *Development* 121, 37–51. 364
- 365 7. Mitsiadis, T. A., Muramatsu, T., Muramatsu,
366 H., and Thesleff, I. (1995) Midkine (MK), a
367 heparin-binding growth/differentiation factor,
368 is regulated by retinoic acid and epithelial-
369 mesenchymal interactions in the developing
370 mouse tooth, and affects cell proliferation and
371 morphogenesis, *J Cell Biol* 129, 267–281.
- 372 8. Mitsiadis, T. A., Dicou, E., Joffre, A., and
373 Magloire, H. (1992) Immunohistochemical
374 localization of nerve growth factor (NGF) and
375 NGF receptor (NGF-R) in the developing first
376 molar tooth of the rat, *Differentiation* 49,
377 47–61.
- 378 9. Mekada, E., Yamaizumi, M., and Okada, Y.
379 (1978) An attempt to separate mononuclear
380 cells fused with human red blood cell-ghosts
381 from a cell mixture treated with HVJ (Sendai
382 virus) using a fluorescence activated cell sorter
383 (FACS II), *J Histochem Cytochem* 26, 62–67.

Author Queries

Chapter No.: 15 0001507499

Queries	Details Required	Author's Response
AU1	Please check whether the edits made to the sentence “The classification of OTs most used...” are ok.	
AU2	Please check whether it should be “pried” instead of “prized” in the sentence “In those OTs that expand the jaws...”.	
AU3	Please check whether it should be “DMEM” instead of “DMEN” in the sentence “The cell suspension is centrifuged...”; comment applicable for all similar usages.	
AU4	Please check whether the edits made to the sentence “These explants are then placed...” are ok.	
AU5	Please check whether “(GFP)” should be deleted from the legend of the Fig. 3.	
AU6	Please check whether the edits made to the sentence “We normally received the tissue...” are ok.	
AU7	Whom does the pronoun “They” implies in the sentence “They select a portion of the tumor...”? Please check and make necessary changes to enhance understand-ability.	